

The system of three vortexes of two dimensional ideal
hydrodynamics as a new example of the (integrable)
Nambu-Poisson mechanics

N.MAKHALDIANI¹,

Joint Institute for Nuclear Research,
Dubna, Moscow region, 141980, Russia

ABSTRACT

A Nambu-Poisson formulation of the system of three ordinary differential equations describing dynamics of three vortexes of the ideal two-dimensional hydrodynamics is given. The system is integrated by quadratures.

¹E-mail: mnv@lcta.jinr.dubna.su

1. The equation of motion of a system of three vortexes can be put in the form [1, 3]

$$\begin{aligned}\dot{M}_1 &= \Gamma_1 M_1 (M_2 - M_3), \\ \dot{M}_2 &= \Gamma_2 M_2 (M_3 - M_1), \\ \dot{M}_3 &= \Gamma_3 M_3 (M_1 - M_2).\end{aligned}\tag{1}$$

Indeed, it is well known [2] that the system of N vortexes can be described by the following system of differential equations:

$$\dot{Z}_n = i \sum_{m \neq n}^N \frac{\Gamma_m}{Z_n^* - Z_m^*}.\tag{2}$$

Then it is easy to verify that the quantities

$$\begin{aligned}M_1 &= |Z_2 - Z_3|^2, \\ M_2 &= |Z_3 - Z_1|^2, \\ M_3 &= |Z_1 - Z_2|^2\end{aligned}\tag{3}$$

satisfy the system (1) after changing the time parameter as follows:

$$dt = \frac{M_1 M_2 M_3}{4S_\Delta} d\tau = (M_1 M_2 M_3)^{\frac{1}{2}} R d\tau,\tag{4}$$

where S_Δ is the area of the triangle with vertexes in the centres of the vortexes and R is the radius of the circle with the vortexes on it.

The system (1) has the integrals of motion

$$\begin{aligned}H_1 &= \sum_{i=1}^3 \frac{M_i}{\Gamma_i}, \\ H_2 &= \sum_{i=1}^3 \frac{\ln M_i}{\Gamma_i}\end{aligned}\tag{5}$$

and can be presented in the form

$$\begin{aligned}\dot{M}_i &= \omega_{ijk} \frac{\partial H_1}{\partial M_j} \frac{\partial H_2}{\partial M_k} \\ &= \{M_i, H_1, H_2\} = \omega_{ijk} \frac{1}{\Gamma_j} \frac{1}{\Gamma_k M_k},\end{aligned}\tag{6}$$

where

$$\begin{aligned}\omega_{ijk} &= \epsilon_{ijk} \rho, \\ \rho &= \Gamma_1 \Gamma_2 \Gamma_3 M_1 M_2 M_3\end{aligned}\tag{7}$$

and the Nambu-Poisson bracket of the functions A, B, C on the three-dimensional phase space M^3 is

$$\{A, B, C\} = \omega_{ijk} \frac{\partial A}{\partial M_i} \frac{\partial B}{\partial M_j} \frac{\partial C}{\partial M_k}.\tag{8}$$

The fundamental bracket is

$$\{M_1, M_2, M_3\} = \omega_{ijk}. \quad (9)$$

Then we can again change the time parameter as

$$d\tau = \rho du \quad (10)$$

and obtain Nambu's mechanics [4]

$$\begin{aligned} \dot{M}_i &= \epsilon_{ijk} \frac{\partial H_1}{\partial M_j} \frac{\partial H_2}{\partial M_k}, \\ \dot{M}_1 &= \frac{M_2 - M_3}{\Gamma_2 \Gamma_3 M_2 M_3}, \\ \dot{M}_2 &= \frac{M_3 - M_1}{\Gamma_3 \Gamma_1 M_3 M_1}, \\ \dot{M}_3 &= \frac{M_1 - M_2}{\Gamma_1 \Gamma_2 M_1 M_2}. \end{aligned} \quad (11)$$

2. The second-order, ternary, Nambu-Poisson structure (6-9) reduces to the two first-order, binary, (Nambu-)Poisson structures[3]

$$\begin{aligned} \{M_i, M_j\}_1 &= (\{M_i, M_j, H_1\} = \omega_{ijk} \frac{\partial H_1}{\partial M_k} = \omega_{ijk} \frac{1}{\Gamma_k}) = \omega_{ij}^1, \\ \{M_i, M_j\}_2 &= (\{M_i, M_j, H_2\} = \omega_{ijk} \frac{\partial H_2}{\partial M_k} = \omega_{ijk} \frac{1}{\Gamma_k M_k}) = \omega_{ij}^2. \end{aligned} \quad (12)$$

These Poisson structures are reducible, there are nontrivial functions H_1 and H_2 for which hold

$$\begin{aligned} \{A, H_1\}_1 &= 0, \\ \{A, H_2\}_2 &= 0 \end{aligned} \quad (13)$$

for any function A .

3. The variables M_n , $n = 1, 2, 3$, are non-negative (semi-bounded), so it is convenient to replace them with free variables x_n

$$x_n = \ln M_n, \quad n = 1, 2, 3. \quad (14)$$

The equation of motion (1), integrals (5) and Nambu-Poisson structures (8-12) take the following form:

$$\begin{aligned} \dot{x}_1 &= \Gamma_1(e^{x_2} - e^{x_3}), \\ \dot{x}_2 &= \Gamma_2(e^{x_3} - e^{x_1}), \\ \dot{x}_3 &= \Gamma_3(e^{x_1} - e^{x_2}), \end{aligned} \quad (15)$$

$$H_1 = \sum_{i=1}^3 \frac{e^{x_i}}{\Gamma_i}, \quad (16)$$

$$H_2 = \sum_{i=1}^3 \frac{x_i}{\Gamma_i},$$

$$\begin{aligned} \dot{x}_i &= \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{\partial H_1}{\partial x_j} \frac{\partial H_2}{\partial x_k} \\ &= \{x_i, H_1, H_2\} = \{x_i, H_1\}_2 = -\{x_i, H_2\}_1, \end{aligned} \quad (17)$$

$$\{A, B, C\} = \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{\partial A}{\partial x_i} \frac{\partial B}{\partial x_j} \frac{\partial C}{\partial x_k}, \quad (18)$$

$$\{A, B\}_1 = \omega_{ij}^1 \frac{\partial A}{\partial x_i} \frac{\partial B}{\partial x_k},$$

$$\{A, B\}_2 = \omega_{ij}^2 \frac{\partial A}{\partial x_i} \frac{\partial B}{\partial x_j},$$

$$\omega_{ij}^1 = \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{\partial H_1}{\partial x_k} \quad (19)$$

$$= \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{1}{\Gamma_k x_k},$$

$$\omega_{ij}^2 = \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{\partial H_2}{\partial x_k}$$

$$= \Gamma_1 \Gamma_2 \Gamma_3 \epsilon_{ijk} \frac{1}{\Gamma_k}.$$

4. For the system of three equations (15) we have two integrals of motion (16), so the system (15) is integrable by quadratures [5, 6]. From H_2 it follows that

$$x_3 = \Gamma_3 \left(H_2 - \frac{x_1}{\Gamma_1} - \frac{x_2}{\Gamma_2} \right). \quad (20)$$

Inserting (20) into the expression of H_1 , we find that

$$\frac{\exp x_2}{\Gamma_2} + \frac{\exp(-\frac{\Gamma_3}{\Gamma_2} x_2)}{\Gamma_3} \exp(\Gamma_3 (H_2 - \frac{x_1}{\Gamma_1})) = H_1 - \frac{\exp x_1}{\Gamma_1}. \quad (21)$$

Now we see that x_2 can be found from (21) as an elementary function of x_1 , when

$$\begin{aligned} \Gamma_3 &= -\Gamma_2, \\ &= -2\Gamma_2, \\ &= -3\Gamma_2, \\ &= -4\Gamma_2, \\ &= \Gamma_2, \\ &= \frac{1}{2}\Gamma_2, \\ &= \frac{1}{3}\Gamma_2. \end{aligned} \quad (22)$$

For the general case, equation (21) defines x_2 as a new transcendental function $n_1(x_1)$. Then the equation for x_1 takes the form

$$\dot{x}_1 = \Gamma_1(e^{n_1(x_1)}(1 + \frac{\Gamma_3}{\Gamma_2}) - \Gamma_3 H_1) + \Gamma_3 e^{x_1} \equiv n_2(x_1) \quad (23)$$

and x_1 is defined by the following quadrature:

$$N(x_1) \equiv \int_{x_{10}}^{x_1} \frac{dx}{n_2(x)} = \tau - \tau_0. \quad (24)$$

Elsewhere we consider general methods of non-linear (Nambu-)Poisson algebras [7] analysis for our model as well as detailed analysis of the formal solution.

It is a pleasure to thank Dr. S.A. Gogilidze and other members of the seminar $\odot NM\pi$ for stimulating discussion.

References

- [1] A.V. Meleshko, N.N. Konstantinov, Dynamics of vortex systems. Kiev: Naukova dumka, 1993.
- [2] H. Aref, Ann.Rev.Fluid Mech.15(1983)345.
- [3] A.V. Borisov and A.V. Cigvincev, Regular & Chaotic Dynamics 1(1996)29.
- [4] Y. Nambu, Phys.Rev.D7(1973)2405.
- [5] W. Gröbli, Spezielle Probleme über die Bewegung geradliniger paralleler Wirbelfäden. Zürich, Zürcher und Furrer, 1877, 86pp.
- [6] E.A. Novikov, Sov. Phys. JETP 41(1975)937.
- [7] N.V. Karasev and V.P. Maslov, Nonlinear Poisson brackets, geometry and quantization. Translation of Math. Monographs, vol. 119, Amer. Math. Soc., Providence, 1993.